

ISBN 82-553-0479-7

Mathematics  
No 4 - March 10

1982

DEFORMATIONS OF REFLEXIVE  
SHEAVES OF RANK 2 ON  $\mathbb{P}^3$

by

Jan Oddvar Kleppe  
Inst. of Math., University of Oslo

DEFORMATIONS OF REFLEXIVE  
SHEAVES OF RANK 2 ON  $\mathbb{P}_k^3$

In this paper we study deformations of reflexive sheaves of rank 2 on  $\mathbb{P} = \mathbb{P}_k^3$  where  $k$  is an algebraically closed field of any characteristic. Let  $\underline{F}$  be a reflexive sheaf with a section  $s \in H^0(\underline{F})$  whose corresponding scheme of zeros is a curve  $C$  in  $\mathbb{P}$ . Moreover let  $M = M(c_1, c_2, c_3)$  be the (coarse) moduli space of stable reflexive sheaves with Chern classes  $c_1, c_2$  and  $c_3$ . The study of how the deformations of  $C \subseteq \mathbb{P}$  correspond to the deformations of the reflexive sheaf  $\underline{F}$  leads to a nice relationship between the local ring  $O_{H,C}$  of the Hilbert scheme  $H = H(d, g)$  of curves of degree  $d$  and arithmetic genus  $g$  at  $C \subseteq \mathbb{P}$  and the corresponding local ring  $O_{M,\underline{F}}$  of  $M$  at  $\underline{F}$ . In this paper we consider some examples where we use this relationship. In particular we prove that the moduli spaces  $M(0, 13, 74)$  and  $M(-1, 14, 88)$  contain generically non-reduced components.

I would like to thank Olav Arnfinn Laudal and Stein Arild Strømme for discussions and comments.

1. Deformations of a reflexive sheaf with a section.

If  $\text{Def}_{\underline{F}}$  is the local deformation functor of  $\underline{F}$  defined on the category  $\underline{1}$  of local artinian  $k$ -algebras with residue field then it is well known that  $\text{Ext}_{O_{\mathbb{P}}}^1(\underline{F}, \underline{F})$  is the tangent space of  $\text{Def}_{\underline{F}}$  and that  $\text{Ext}_{O_{\mathbb{P}}}^2(\underline{F}, \underline{F})$  contains the obstructions of deformation. See [H3]. To deform the pair  $(\underline{F}, s)$  we consider the functor

$$\text{Def}_{\underline{F}, s} : \underline{1} \rightarrow \underline{\text{Sets}}$$

defined by

$$\text{Def}_{\underline{F},s}(R) = \{O_{\mathbb{P}_R} \xrightarrow{s_R} \underline{F}_R \mid \underline{F}_R \in \text{Def}_{\underline{F}}(R) \text{ and } s_R \otimes_R 1_k = s\} / \sim$$

where  $\mathbb{P}_R = \mathbb{P} \times \text{Spec}(R)$  and where  $1_k : k \rightarrow k$  is the identity. Two deformations  $(\underline{F}_R, s_R)$  and  $(\underline{F}'_R, s'_R)$  are equivalent if there exist isomorphisms  $O_{\mathbb{P}_R} \xrightarrow{\sim} O_{\mathbb{P}_R}$ ,  $\underline{F}_R \xrightarrow{\sim} \underline{F}'_R$  and a commutative diagram

$$\begin{array}{ccc} O_{\mathbb{P}_R} & \xrightarrow{s_R} & \underline{F}_R \\ \cong \downarrow & \circ & \downarrow \cong \\ O_{\mathbb{P}_R} & \xrightarrow{s'_R} & \underline{F}'_R \end{array}$$

such that  $s_R \otimes_R 1_k = s'_R \otimes_R 1_k$ . In fact we also identify the given pair  $(\underline{F}, s)$  with any  $(\underline{F}', s')$  where  $s' \in H^0(\mathbb{P}, \underline{F}')$  if they fit together into such a commutative diagram.

Proposition 1.1. (i) The tangent space of  $\text{Def}_{\underline{F},s}$  is

$$\text{Ext}_{O_{\mathbb{P}}}^1(\underline{I}_C(c_1), \underline{F}) \text{ where } \underline{I}_C = \ker(O_{\mathbb{P}} \rightarrow O_C), \text{ and}$$

$$\text{Ext}_{O_{\mathbb{P}}}^2(\underline{I}_C(c_1), \underline{F}) \text{ contains the obstructions of deformations.}$$

(ii) The natural

$$\varphi : \text{Def}_{\underline{F},s} \rightarrow \text{Def}_{\underline{F}}$$

is a smooth morphism of functors on  $\underline{1}$  provided

$$H^1(\underline{F}) = 0$$

By the correspondence [H3, 4.1] there is a curve  $C = (s)_0 \subseteq \mathbb{P}$  and an exact sequence

$$\xi : 0 \rightarrow O_{\mathbb{P}} \xrightarrow{s} \underline{F} \rightarrow \underline{I}_C(c_1) \rightarrow 0$$

associated to  $(\underline{F}, s)$ . The condition  $H^1(\underline{F}) = 0$  is therefore equivalent to

$$H^1(\underline{I}_C(c_1)) = 0$$

Proof of (i). Using [L2, §2] or [K1, 1.2] we know that there is a spectral sequence

$$E_2^{p,q} = \lim_{\leftarrow} (p) \left\{ \begin{array}{ccc} \text{Ext}^q(\underline{F}, \underline{F}) & & \text{Ext}^q(\mathcal{O}_{\mathbb{P}}, \mathcal{O}_{\mathbb{P}}) \\ & \searrow \alpha^q & \swarrow \\ & \text{Ext}^q(\mathcal{O}_{\mathbb{P}}, \underline{F}) & \end{array} \right\}$$

converging to some group  $A^{(\cdot)}$  where  $A^1$  is the tangent space of  $\text{Def}_{\underline{F}, s}$  and  $A^2$  contains the obstructions of deformation. Since  $E_2^{p,q} = 0$  for  $p \geq 2$ , we have an exact sequence

$$0 \rightarrow E_2^{1, q-1} \rightarrow A^q \rightarrow E_2^{0, q} \rightarrow 0$$

Moreover

$$\text{Ext}^q(\mathcal{O}_{\mathbb{P}}, \mathcal{O}_{\mathbb{P}}) = 0 \text{ for } q > 0 \text{ and } \text{Ext}^q(\mathcal{O}_{\mathbb{P}}, \underline{F}) = H^q(\underline{F}) \text{ for any } q,$$

and this gives

$$E_2^{0, q} = \ker \alpha^q \text{ and } E_2^{1, q} = \text{coker } \alpha^q \text{ for } q > 0.$$

Observe also that

$$E_2^{1, 0} = \lim_{\leftarrow} (1) \left\{ \begin{array}{ccc} \text{Hom}(\underline{F}, \underline{F}) & & \text{Hom}(\mathcal{O}_{\mathbb{P}}, \mathcal{O}_{\mathbb{P}}) \\ & \searrow \alpha^0 & \swarrow \\ & \text{Hom}(\mathcal{O}_{\mathbb{P}}, \underline{F}) & \end{array} \right\} = \text{coker } \alpha^0$$

because  $\text{Hom}(\mathcal{O}_{\mathbb{P}}, \mathcal{O}_{\mathbb{P}}) \subseteq \text{Hom}(\underline{F}, \underline{F})$ . We therefore have an exact sequence

$$0 \rightarrow \text{coker } \alpha^{q-1} \rightarrow A^q \rightarrow \ker \alpha^q \rightarrow 0$$

for any  $q > 0$ . Combining with the long exact sequence

$$\begin{aligned}
 (*) \quad & \longrightarrow \text{Hom}(\underline{F}, \underline{F}) \xrightarrow{\alpha^0} H^0(\underline{F}) \longrightarrow \text{Ext}^1(\underline{L}_C(c_1), \underline{F}) \xrightarrow{\varphi^1} \text{Ext}^1(\underline{F}, \underline{F}) \\
 & \xrightarrow{\alpha^1} H^1(\underline{F}) \longrightarrow \text{Ext}^2(\underline{L}_C(c_1), \underline{F}) \xrightarrow{\varphi^2} \text{Ext}^2(\underline{F}, \underline{F}) \xrightarrow{\alpha^2} H^2(\underline{F}) \longrightarrow
 \end{aligned}$$

deduced from the short exact sequence

$$0 \rightarrow 0_{\mathbb{P}} \xrightarrow{s} \underline{F} \rightarrow \underline{L}_C(c_1) \rightarrow 0,$$

we find isomorphisms

$$A^q \simeq \text{Ext}^q(\underline{L}_C(c_1), \underline{F}) \quad \text{for } q > 0.$$

(ii) Let  $S \rightarrow R$  be a morphism in  $\underline{\mathcal{L}}$  whose kernel  $\mathcal{C}$  is a  $k$ -module via  $R \twoheadrightarrow k$ , let  $s_R : 0_{\mathbb{P}_R} \rightarrow \underline{F}_R$  be a deformation of  $s : 0_{\mathbb{P}} \rightarrow \underline{F}$  to  $R$ , and let  $\underline{F}_S$  be a deformation of  $\underline{F}_R$  to  $S$ . To prove the smoothness of  $\varphi$ , we must find a morphism  $s_S$ ,

$$s_S : 0_{\mathbb{P}_S} \rightarrow \underline{F}_S$$

such that  $s_S \otimes_S 1_R = s_R$ , i.e. we must prove that  $s_R \in H^0(\underline{F}_R)$  is contained in the image of  $H^0(\underline{F}_S) \rightarrow H^0(\underline{F}_R)$ . Since

$$0 \rightarrow \underline{F} \otimes_k \mathcal{C} \rightarrow \underline{F}_S \rightarrow \underline{F}_R \rightarrow 0$$

is exact and since  $H^1(\underline{F}) = 0$  by assumption, we see that  $H^0(\underline{F}_S) \rightarrow H^0(\underline{F}_R)$  is surjective and we are done.

Remark 1.2. In the exact sequence (\*) of this proof,  $\varphi^1$  is the tangent map of  $\varphi : \text{Def}_{\underline{F}, s} \rightarrow \text{Def}_{\underline{F}}$  and  $\varphi^2$  maps "obstructions to obstructions". In fact  $\varphi$  is a morphism of principal homogeneous spaces via  $\varphi^1$ . Using this it is in general rather easy to prove the smoothness of  $\varphi$  directly from the surjectivity of  $\varphi^1$  and the injectivity of  $\varphi^2$ . This gives another proof of (1.1. ii).

2. The relationship between the deformations of a reflexive sheaf with a section and the deformations of the corresponding curve.

Let  $\underline{F}$ ,  $s \in H^0(\underline{F})$  and  $\underline{I} = \underline{I}_C = \ker(O_{\mathbb{P}} \rightarrow O_C)$  be as in the preceding section, and let  $\text{Def}_{\underline{I}} : \underline{1} \rightarrow \text{Sets}$  be the deformation functor of the  $O_{\mathbb{P}}$ -Module  $\underline{I}$ . Then there is a natural map

$$\psi : \text{Def}_{\underline{F}, s} \rightarrow \text{Def}_{\underline{I}}$$

defined by

$$\psi(\underline{F}_R, s_R) = \underline{M}_R \otimes (O_{\mathbb{P}}(-c_1) \otimes_k R)$$

where  $\underline{M}_R = \text{coker } s_R$ . If  $\text{Hilb}_C : \underline{1} \rightarrow \text{Sets}$  is the local Hilbert functor at  $C \subseteq \mathbb{P}$ , we have also a natural map

$$\text{Hilb}_C \rightarrow \text{Def}_{\underline{I}}$$

of functors on  $\underline{1}$ . Recall that  $C$  is locally Cohen Macaulay and equidimensional [H3, 4.1].

Proposition 2.1. (i) The natural morphism

$$\text{Hilb}_C \rightarrow \text{Def}_{\underline{I}}$$

is an isomorphism of functors.

(ii) If  $H^1(\underline{F}(-4)) = 0$ , then

$$\psi : \text{Def}_{\underline{F}, s} \rightarrow \text{Def}_{\underline{I}}$$

is a smooth morphism of functors on  $\underline{1}$ .

Observe also that

$$H^1(\underline{F}(-4)) \simeq H^1(\underline{I}_C(c_1-4))$$

and moreover by duality that

$$\text{Ext}_{O_{\mathbb{P}}}^2(\underline{I}_C(c_1), O_{\mathbb{P}}) = H^1(\underline{I}_C(c_1-4))^{\vee}.$$

Proof of (i) If  $\underline{N}_C = \underline{\text{Hom}}_{O_{\mathbb{P}}}(\underline{I}, O_C)$  is the normal bundle of  $C$  in  $\mathbb{P}$ , we proved in [K1, 2.2] that

$$H^i(\underline{N}_C) \simeq \text{Ext}_{O_{\mathbb{P}}}^{i+1}(\underline{I}, \underline{I}) \quad \text{for } i = 0, 1$$

as a consequence of the fact that the projective dimension of the  $O_{\mathbb{P}}$ -Module  $\underline{I}$  is 1, from which the conclusion of (i) is easy to understand. We will, however, give a direct proof.

To construct the inverse of  $\text{Hilb}_C(R) \rightarrow \text{Def}_{\underline{I}}(R)$ , let  $\underline{M}_R$  be a deformation of  $\underline{I}$  to  $R$ . Observe that there is an exact sequence

$$(*) \quad 0 \rightarrow \underline{E} \rightarrow \bigoplus_{i=1}^{r+1} O_{\mathbb{P}}(-n_i) \xrightarrow{f} \underline{I} \rightarrow 0$$

where  $\underline{E}$  is a vector bundle on  $\mathbb{P}$  of rank  $r$ .  $\wedge^r \underline{E}$  is therefore invertible, and we can identify it with  $O_{\mathbb{P}}(d_1)$  where  $d_1 = -\sum n_i$ . If  $\underline{P} = \bigoplus O_{\mathbb{P}}(-n_i)$ , then there is a complex

$$(**) \quad \underline{E} \rightarrow \underline{P} \xrightarrow{\sim} (\wedge^r \underline{P})^{\vee}(d_1) \rightarrow (\wedge^r \underline{E})^{\vee}(d_1) = O_{\mathbb{P}}$$

and it is well known that the maps  $\underline{P} \xrightarrow{f} \underline{I} \subseteq O_{\mathbb{P}}$  and  $\underline{P} \rightarrow O_{\mathbb{P}}$  deduced from (\*) and (\*\*) respectively are equal up to a unit of  $k$ . We can assume equality. Now since  $\underline{M}_R$  is a lifting of  $\underline{I}$  to  $R$ , there is a map

$$f_R : \underline{P}_R = \bigoplus_{i=1}^{r+1} O_{\mathbb{P}_R}(-n_i) \rightarrow \underline{M}_R$$

such that  $f_R \otimes_R 1_k = f : \underline{P} \rightarrow \underline{I}$ . By Nakayama's lemma,  $f_R$  is surjective. Moreover if  $\underline{E}_R = \ker f_R$ , we easily see that  $\underline{E}_R \otimes_R k$

and  $\underline{E}_R$  is  $R$ -flat. It follows that  $\underline{E}_R$  is a locally free  $\mathcal{O}_{\mathbb{P}^r}$ -Module of rank  $r$  satisfying

$$\wedge^r \underline{E}_R = \mathcal{O}_{\mathbb{P}^r}(d_1).$$

Furthermore there is a complex

$$\underline{E}_R \rightarrow \underline{P}_R \simeq (\wedge^r \underline{P}_R)^\vee(d_1) \rightarrow (\wedge^r \underline{E}_R)^\vee(d_1) = \mathcal{O}_{\mathbb{P}^r}$$

which proves the existence of an  $\mathcal{O}_{\mathbb{P}^r}$ -linear map

$$\alpha : \underline{M}_R \rightarrow \mathcal{O}_{\mathbb{P}^r}$$

which reduces to the natural inclusion  $\underline{I} \subseteq \mathcal{O}_{\mathbb{P}^r}$  via  $(-)^{\otimes_R k}$ . It

is easy to see that  $\alpha$  is injective, that  $\text{coker } \alpha$  is  $R$ -flat

and that  $\text{coker } \alpha^{\otimes_R k} = \mathcal{O}_C$ . We therefore have a deformation

$C_R \subseteq \mathbb{P}^r$  of  $C \subseteq \mathbb{P}^r$ . Finally to see that the inverse

of  $\text{Hilb}_C(R) \rightarrow \text{Def}_{\underline{I}}(R)$  is well-defined, let  $\beta : \underline{M}'_R \xrightarrow{\simeq} \underline{M}_R$  and

$\alpha' : \underline{M}'_R \rightarrow \mathcal{O}_{\mathbb{P}^r}$  be  $\mathcal{O}_{\mathbb{P}^r}$ -linear maps such that  $\beta^{\otimes_R 1_k}$  is the identity on  $\underline{I}$  and  $\alpha'^{\otimes_R 1_k}$  is the natural inclusion  $\underline{I} \subseteq R$ .

(We do not assume  $\alpha'\beta = \alpha$ ). We claim that  $\text{Im } \alpha' = \text{Im } \alpha$ . In fact since

$$\text{Ext}_{\mathcal{O}_{\mathbb{P}^r}}^i(\mathcal{O}_C, \mathcal{O}_{\mathbb{P}^r}) = 0 \quad \text{for } i = 0, 1,$$

we have

$$k = \text{Hom}_{\mathcal{O}_{\mathbb{P}^r}}(\mathcal{O}_{\mathbb{P}^r}, \mathcal{O}_{\mathbb{P}^r}) \xrightarrow{\simeq} \text{Hom}_{\mathcal{O}_{\mathbb{P}^r}}(\underline{I}, \mathcal{O}_{\mathbb{P}^r}).$$

We deduce that the map

$$R = \text{Hom}_{\mathcal{O}_{\mathbb{P}^r}}(\mathcal{O}_{\mathbb{P}^r}, \mathcal{O}_{\mathbb{P}^r}) \rightarrow \text{Hom}_{\mathcal{O}_{\mathbb{P}^r}}(\underline{M}_R, \mathcal{O}_{\mathbb{P}^r})$$

induced by  $\alpha$ , is surjective. Hence

$$\alpha'\beta = \alpha.$$



for some  $r \in R$ , and since  $\alpha' \beta \otimes 1_k = \alpha \otimes 1_k$  is the natural inclusion  $\underline{I} \subseteq \underline{O}_{\mathbb{P}}$ ,  $r$  is a unit and we are done.

(ii) Let  $S \rightarrow R$ ,  $\mathcal{O}$  and  $s_R : \underline{O}_{\mathbb{P}_R} \rightarrow \underline{F}_R$  be as in the proof of (1.1 ii). Moreover let  $\underline{M}_R = \text{coker } s_R$ , and let  $\underline{M}_S$  be a deformation of  $\underline{M}_R$  to  $S$ . To prove smoothness we must find a deformation

$$s_S : \underline{O}_{\mathbb{P}_S} \rightarrow \underline{F}_S$$

with cokernel  $\underline{M}_S$  such that  $s_S \otimes_S 1_R = s_R$ . By theory of extensions it is sufficient to prove that the map

$$\text{Ext}_{\underline{O}_{\mathbb{P}_S}}^1(\underline{M}_S, \underline{C}_{\mathbb{P}_S}) \rightarrow \text{Ext}_{\underline{O}_{\mathbb{P}_R}}^1(\underline{M}_R, \underline{O}_{\mathbb{P}_R})$$

induced by  $(-)\otimes_S R$  is surjective. Modulo isomorphisms we refind this map in the long exact sequence

$$\rightarrow \text{Ext}_{\underline{O}_{\mathbb{P}_S}}^1(\underline{M}_S, \underline{O}_{\mathbb{P}_S} \otimes \mathcal{O}) \rightarrow \text{Ext}_{\underline{O}_{\mathbb{P}_S}}^1(\underline{M}_S, \underline{O}_{\mathbb{P}_S}) \rightarrow \text{Ext}_{\underline{O}_{\mathbb{P}_R}}^1(\underline{M}_S, \underline{O}_{\mathbb{P}_R}) \rightarrow \text{Ext}_{\underline{O}_{\mathbb{P}_S}}^2(\underline{M}_S, \underline{O}_{\mathbb{P}_S} \otimes \mathcal{O}).$$

Since  $\text{Ext}_{\underline{O}_{\mathbb{P}_S}}^2(\underline{M}_S, \underline{O}_{\mathbb{P}_S} \otimes_S \mathcal{O}) \simeq \text{Ext}_{\underline{O}_{\mathbb{P}}}^2(\underline{I}_C(c_1), \underline{O}_{\mathbb{P}}) \otimes \mathcal{O} = 0$  by

assumption, we are done.

Remark 2.2. The short exact sequence

$$\xi : 0 \rightarrow \underline{O}_{\mathbb{P}} \xrightarrow{s} \underline{F} \rightarrow \underline{I}_C(c_1) \rightarrow 0$$

induces a long exact sequence

$$\begin{aligned} \rightarrow \text{Ext}_{\underline{O}_{\mathbb{P}}}^1(\underline{I}_C(c_1), \underline{O}_{\mathbb{P}}) \rightarrow \text{Ext}_{\underline{O}_{\mathbb{P}}}^1(\underline{I}_C(c_1), \underline{F}) \xrightarrow{\psi^1} \text{Ext}_{\underline{O}_{\mathbb{P}}}^1(\underline{I}_C, \underline{I}_C) \rightarrow \\ \text{Ext}_{\underline{O}_{\mathbb{P}}}^2(\underline{I}_C(c_1), \underline{O}_{\mathbb{P}}) \rightarrow \text{Ext}_{\underline{O}_{\mathbb{P}}}^2(\underline{I}_C(c_1), \underline{F}) \xrightarrow{\psi^2} \text{Ext}_{\underline{O}_{\mathbb{P}}}^2(\underline{I}_C, \underline{I}_C) \rightarrow \end{aligned}$$

where  $\psi^1$  is the tangent map of  $\psi$  or more generally,  $\psi$  is a map of principal homogeneous spaces via  $\psi^1$  and  $\psi^2$  maps "obstructions to obstructions". As remarked in (1.2), the smoothness of  $\psi$  follows therefore from the surjectivity of  $\psi^1$  and the injectivity of  $\psi^2$ .

Remark 2.3. Let  $\xi$  be the extension

$$0 \rightarrow 0_{\mathbb{P}} \xrightarrow{s} \underline{F} \rightarrow \underline{I}_C(c_1) \rightarrow 0$$

and let  $\text{Def}_{C, \xi} : \underline{1} \rightarrow \underline{\text{Sets}}$  be the functor defined by

$$\text{Def}_{C, \xi}(R) = \left\{ (C_R, \xi_R) \left| \begin{array}{l} (C_R \subseteq \mathbb{P}_R) \in \text{Hilb}_C(R) \text{ and } \xi_R \in \\ \text{Ext}^1(\underline{I}_{C_R}(c_1), 0_{\mathbb{P}_R}) \text{ satisfies} \\ \xi_R \otimes_R k = \xi \end{array} \right. \right\} / \sim$$

Two deformations  $(C_R, \xi_R)$  and  $(C'_R, \xi'_R)$  are equivalent if  $C_R = C'_R \subseteq \mathbb{P}_R$  and if there is a commutative diagram

$$\begin{array}{ccccccc} \xi'_R : 0 & \rightarrow & 0_{\mathbb{P}_R} & \rightarrow & \underline{F}_R & \rightarrow & \underline{I}_{C_R}(c_1) \rightarrow 0 \\ & & \downarrow & \cdot & \downarrow & \cdot & \parallel \quad 1 \\ \xi_R : 0 & \rightarrow & 0_{\mathbb{P}_R} & \rightarrow & \underline{F}_R & \rightarrow & \underline{I}_{C_R}(c_1) \rightarrow 0, \end{array}$$

both reducing to the extension  $\xi$  via  $(-)^{\otimes_R k}$ . In the same way we identify the given  $(C, \xi)$  with any  $(C', \xi')$  provided  $C = C'$  and  $\xi' = u\xi$  for some unit  $u \in k^*$ . Note that we may in this definition of equivalence replace the identity 1 on  $\underline{I}_{C_R}(c_1)$  by any  $0_{\mathbb{P}_R}$  linear map. See [Ma 2, 6.1] and recall  $\text{Hom}(\underline{I}_C, \underline{I}_C) = k$ . Now there is a forgetful map

$$c : \text{Def}_{C, \xi} \rightarrow \text{Def}_{\underline{F}, s},$$

and using (2.1i) we immediately have an inverse of  $\alpha$ .

Hence  $\alpha$  is an isomorphism. Observe that we might construct the inverse of  $\alpha(R)$  for  $R \in \text{ob } \underline{1}$  by considering the invertible sheaf  $\det \underline{F}_R$  on  $\mathbb{P}_R$ . See [Ma 1, 4.2] or [G, 4.1]. In fact if  $(\underline{F}_R, s_R)$  is given, there is an  $\mathbb{P}_R$  a morphism

$$i : \wedge^2 \underline{F}_R \rightarrow \det \underline{F}_R \simeq \mathcal{O}_{\mathbb{P}_R}(c_1)$$

and a complex

$$0 \rightarrow \mathcal{O}_{\mathbb{P}_R} \xrightarrow{s_R} \underline{F}_R \xrightarrow{i[(-) \wedge s_R]} \mathcal{O}_{\mathbb{P}_R}(c_1)$$

which after the tensorization  $(-)^{\otimes_R k}$  is exact. Hence

$$0 \rightarrow \mathcal{O}_{\mathbb{P}_R} \xrightarrow{s_R} \underline{F}_R \rightarrow \text{coker } s_R \rightarrow 0$$

is exact,  $\text{coker } s_R$  is  $R$ -flat and  $\text{coker } s_R \simeq \mathcal{O}_{\mathbb{P}_R}(c_1)$ , and putting this together, we can find an inverse of  $\alpha(R)$ .

One should compare the isomorphism of  $\alpha$  with [H 3, 4.1] which implies that there is a bijection between the set of pairs  $(\underline{F}, s)$  and the set of  $(C, \xi)$  moduli equivalence under certain conditions on the pairs. Thinking of these families of pairs as moduli spaces, [H 3, 4.1] establishes a bijection on the  $k$ -points of these spaces while the isomorphism of  $\alpha$  takes care of the scheme structure as well.

To be more precise we claim that there is a quasiprojective scheme  $D$  parametrizing equivalent pairs  $(C, \xi)$  where

- 1)  $C$  is an equidimensional Cohen Macaulay curve and where
- 2) the extension  $\xi : 0 \rightarrow \mathcal{O}_{\mathbb{P}} \rightarrow \underline{F} \rightarrow \underline{I}_C(c_1) \rightarrow 0$  is such that  $\underline{F}$  is a stable reflexive sheaf.

Moreover there are projection morphisms

$$\begin{array}{ccc}
 & D & \xrightarrow{q} & H(d,g) \\
 (*) & p \downarrow & & \\
 & M(c_1, c_2, c_3) & & 
 \end{array}$$

defined by  $p(\underline{F}_K, s_K) = \underline{F}_K$  and  $q(C_K, \xi_K) = C_K$  for a geometric  $K$ -point  $(C_K, \xi_K)$  corresponding to  $(\underline{F}_K, s_K)$ , such that the fibers of  $p$  and  $q$  are smooth connected schemes. Furthermore,  $p$  is smooth at  $(\underline{F}_K, s_K)$  provided  $H^1(\underline{F}_K) = 0$ , and  $q$  is smooth at  $(C_K, \xi_K)$  provided  $H^1(\underline{I}_{C_K}(c_1-4)) = 0$ .

To indicate why <sup>1)</sup> let  $\underline{Sch}/k$  be the category of locally noetherian  $k$ -schemes and let  $\underline{D} : \underline{Sch}/k \rightarrow \underline{Sets}$  be the functor defined by

$$\underline{D}(S) = \left\{ (C_S, \underline{L}_S, \xi_S) \left| \begin{array}{l} C_S \in \underline{H}(d,g)(S), \underline{L}_S \text{ is invertible on } S \text{ and} \\ \xi_S \in \text{Ext}^1(\underline{I}_{C_S}(c_1), \mathcal{O}_{\mathbb{P} \times S} \otimes \underline{L}_S) \text{ such that} \\ C_S \times_S \text{Spec}(K) \text{ satisfies (1) and } \xi_S \otimes K \neq 0 \\ \text{for any geometric } K\text{-point of } S \end{array} \right. \right\}$$

Two deformations  $(C_S, \underline{L}_S, \xi_S)$  and  $(C'_S, \underline{L}'_S, \xi'_S)$  are equivalent if  $C_S = C'_S$  and if there is an isomorphism  $\tau : \underline{L}_S \rightarrow \underline{L}'_S$  whose induced morphism  $\text{Ext}^1(\underline{I}_{C_S}(c_1), \tau)$  maps  $\xi_S$  onto  $\xi'_S$ . Now if  $U \subseteq H(d,g)$  is the open set of equidimensional Cohen Macaulay curves and if  $C_U \subseteq \mathbb{P} \times U \xrightarrow{\pi} U$  is the restricting of the universal curve to  $U$ , one may prove that  $\underline{E} = \underline{\text{Ext}}^1(\underline{I}_{C_U}(c_1), \mathcal{O}_{\mathbb{P} \times U})$  is a coherent  $\mathcal{O}_{\mathbb{P} \times U}$ -Module, flat over  $U$ . By [EGA, III, 7.7.6] there is a unique coherent  $\mathcal{O}_U$ -Module  $\underline{Q}$  such that

1) For good ideas of this construction, see the appendix [E,S], some of which appears in [S,M,S].

$$\underline{\text{Hom}}_{\mathcal{O}_U}(\underline{Q}, \underline{R}) \simeq \pi_*(\underline{E} \otimes \underline{R})$$

for any quasicoherent  $\mathcal{O}_U$ -Module  $\underline{R}$ . If  $\mathbb{P}(\underline{Q}) = \text{Proj}(\text{Sym}(\underline{Q}))$  is the projective fiber over  $U$  defined by  $\underline{Q}$ , we can use [EGA II, 4.2.3] to prove that

$$\underline{D}(-) \simeq \text{Mor}_k(-, \mathbb{P}(\underline{Q})).$$

Now let  $D \subseteq \mathbb{P}(\underline{Q})$  be the open set whose  $k$ -points are  $(C, \xi)$ ,  $\xi: 0 \rightarrow \mathcal{O}_{\mathbb{P}} \rightarrow \underline{F} \rightarrow \underline{I}_C(c_1) \rightarrow 0$ , where  $\underline{F}$  is a stable reflexive sheaf. Then we have a diagram (\*) where the existence of the morphism  $p$  follows from the definition [Ma 1, 5.5] of the moduli space  $M = M(c_1, c_2, c_3)$ . Moreover since  $\mathbb{P}(\underline{Q})$  represents the functor  $\underline{D}$ , the fiber of  $q: D \rightarrow H(d, g)$  at a  $K$ -point  $C_K \subseteq \mathbb{P}_K$  of  $H(d, g)$  is just  $D \cap \mathbb{P}(\text{Ext}^1(\underline{I}_{C_K}(c_1), \mathcal{O}_{\mathbb{P}_K})^\vee)$  where  $(-)^\vee = \text{Hom}_K(-, K)$ . Moreover if we think of the fiber of  $p$  at a geometric  $K$ -point  $\underline{F}_K$  of  $M$  as those sections  $s \in H^0(\underline{F}_K)$  where  $(s)_0$  is a curve, we understand that the fiber is an open subscheme of the linear space  $\mathbb{P}(H^0(\underline{F}_K)^\vee)$ . In particular the geometric fibers of  $p$  and  $q$  are smooth and connected.

Finally the smoothness of  $p$  and  $q$  at  $(C, \xi)$  follows from (1.1 ii) and (2.1 ii) provided we know that the morphism  $p^*: \mathcal{O}_{M, \underline{F}} \rightarrow \mathcal{O}_{D, (\underline{F}, s)}$  induced by  $p: D \rightarrow M$  makes a commutative diagram

$$\begin{array}{ccc} \text{Def}_{\underline{F}, s} & \simeq & \text{Mor}(\hat{\mathcal{O}}_{D, (\underline{F}, s)}, -) \\ \varphi \downarrow & \circ & \downarrow \text{Mor}(p^*, -) \\ \text{Def}_{\underline{F}} & \simeq & \text{Mor}(\hat{\mathcal{O}}_{M, \underline{F}}, -) \end{array}$$

of horizontal isomorphisms on  $\underline{1}$ . In fact the commutativity from

the definition of a moduli space [Ma 1, 5.5] while the construction of  $M$  implies the lower horizontal isomorphism. See [Ma 2, 6.4] from which we immediately have that the morphism  $\text{Def}_{\underline{F}} \rightarrow \text{Mor}(\hat{O}_{M, \underline{F}}, -)$  is smooth, and since the morphism induces an isomorphism of tangent spaces, both isomorphic to  $\text{Ext}^1(\underline{F}, \underline{F})$ , it must be an isomorphism.

Remark 2.4. In particular the smoothness of  $\text{Def}_{\underline{F}} \rightarrow \text{Mor}(\hat{O}_{M, \underline{F}}, -)$  which is a consequence of the smoothness of the morphism treated in [Ma 2, 6.4], implies that  $O_{M, \underline{F}}$  is a regular local ring if and only if  $\text{Def}_{\underline{F}}$  is a smooth functor on  $\underline{1}$ .

### 3. Non-reduced components of the moduli scheme $M(c_1, c_2, c_3)$ .

One knows that the Hilbert scheme  $H(d, g)$  is not always reduced. In fact if  $g$  is the largest number satisfying  $g \leq \frac{d^2 - 4}{8}$ , we proved in [K1, 3.2.10] that  $H(d, g)$  is non-reduced for every  $d \geq 14$ , and we explicitly described a non-reduced component in terms of the Picard group of a smooth general cubic surface.

Example 3.1. (Mumford [M1]). For  $d = 14$ , we have

$g = \frac{d^2 - 4}{8} = 24$ , and there is an open irreducible subscheme  $U \subseteq H(14, 24)$  of smooth connected curves whose closure  $\bar{U} = W$  makes a non-reduced component, such that for any  $(C \subseteq \mathbb{P}) \in U$ ,

$$h^0(\underline{I}_C(v)) = \begin{cases} 0 & \text{for } v \leq 2 \\ 1 & \text{for } v = 3 \end{cases}$$

$$h^1(\underline{I}_C(v)) = 0 \quad \text{for } v \notin \{3, 4, 5\},$$

$$h^1(O_C(v)) = \begin{cases} 0 & \text{for } v \geq 4 \\ 1 & \text{for } v = 3. \end{cases}$$

See [K1,(3.2.4) and (3.1.3)]. In fact with  $C \subseteq \mathbb{P}$  in  $U$ , there is a global complete intersection of two surfaces of degree 3 and 6 whose corresponding linked curve is a disjoint union of two coniques.

Now let  $C \subseteq \mathbb{P}$  be a smooth connected curve satisfying

$$(*) \quad H^1(\underline{I}_C(c_1)) = 0, \quad H^1(\underline{I}_C(c_1-4)) = 0 \quad \text{and} \quad H^1(\mathcal{O}_C(c_1-4)) \neq 0$$

for some integer  $c_1$ , let  $\xi \in H^0(\omega_C(4-c_1)) = \text{Ext}^1(\underline{I}_C(c_1), \mathcal{O}_{\mathbb{P}})$  be non-trivial, and let  $(\underline{F}, s)$ ,  $s \in H^0(\underline{F})$ , correspond to  $(C, \xi)$  via the usual correspondence. Then  $\underline{F}$  is reflexive, and it is stable (resp. semistable) if and only if  $c_1 > 0$  (resp.  $c_1 \geq 0$ ) and  $C$  is not contained in any surface of degree  $\leq \frac{1}{2}c_1$  (resp.  $< \frac{1}{2}c_1$ ). See [H3, 4.2]. Combining (1.1) and (2.1) with (2.4) in case  $\underline{F}$  is stable, we find that  $\mathcal{O}_{M, \underline{F}}$  is non-reduced iff  $\mathcal{O}_{H, C}$  is non-reduced.

Example 3.2. Let  $(C \subseteq \mathbb{P}) \in H(14, 24)$  belong to the set  $U$  of (3.1) and let  $c_1$  be an integer satisfying (\*), i.e.  $c_1 \leq 2$  or  $c_1 = 6$ .

(i) Let  $c_1 = 6$ . By virtue of (1.1) and (2.1) the hull of  $\text{Def}_{\underline{F}}$  is non-reduced. Moreover  $\underline{F}$  is semistable with Chern classes  $(c_1, c_2, c_3) = (6, 14, 18)$ , and the normalized sheaf  $\underline{F}(-3)$  has Chern classes  $(c'_1, c'_2, c'_3) = (0, 5, 18)$ .

(ii) Let  $c_1 = 2$ . The corresponding reflexive sheaf is stable and must belong to at least one non-reduced component of  $M(2, 14, 74)$ , i.e. of  $M(0, 13, 74)$ .

(iii) With  $c_1 = 1$  we find at least one non-reduced component of  $M(1, 14, 88) \simeq M(-1, 14, 88)$ .

Combining the discussion after (2.3) and in particular the irreducibility of the morphism  $q$  with the irreducibility of the set  $U$  of (3.1), we see that we obtain precisely one non-reduced component of  $M(0,13,74)$  and  $M(-1,14,88)$  in this way.

We will give one more example of a non-reduced component and include a discussion to better understand (1.1) and (2.1). In fact recall [K1,2.3.6] that if an equidimensional Cohen Macaulay curve  $(C \subseteq \mathbb{P}) \in H(d,g)$  is contained in a complete intersection  $V(\underline{F}_1, \underline{F}_2)$  of two surfaces of degree  $f_1 = \deg F_1$  and  $f_2 = \deg F_2$  with

$$H^1(\underline{I}_C(f_i)) = 0 \quad \text{and} \quad H^1(\underline{I}_C(f_i-4)) = 0$$

for  $i = 1, 2$ , and if  $(C' \subseteq \mathbb{P}) \in H' = H(d', g')$  is the linked curve, then  $O_{H,C}$  is reduced iff  $O_{H',C'}$  is reduced. Since any curve  $(C \subseteq \mathbb{P}) \in U$  of (3.1) is contained in a complete intersection  $V(\underline{F}_1, \underline{F}_2)$  of two surfaces of degree  $f_1 = f_2 = 6$ , the linked curves  $C' \subseteq \mathbb{P}$  must belong to at least one (and one may prove to exactly one) non-reduced component<sup>1)</sup>  $W \subseteq H(22, 56)$  of dimension 88. See [K1,2.3.9]. One may see that  $W$  contains smooth connected curves. Moreover using the fact that  $\omega_C(4-f_1-f_2)$  and  $\omega_{C'}(4-f_1-f_2)$  are the sheaves of ideals which define the closed subschemes  $C' \subseteq V(\underline{F}_1, \underline{F}_2)$  and  $C \subseteq V(\underline{F}_1, \underline{F}_2)$  respectively, one proves easily that

$H^0(\underline{I}_C(4)) = 0$ ,  $H^1(\underline{I}_C(v)) = 0$  for  $v \notin \{3, 4, 5\}$  and  $H^1(O_{C'}(5)) \neq 0$   
See [S,P] and [K1,2.3.3].

---

1) The condition  $H^1(\underline{I}_C(f_i-4)) = 0$  implies also that the linked curves  $C' \subseteq \mathbb{P}$  form an open subset of  $H'$ .



Example 3.3. Let  $(C' \subseteq \mathbb{P}) \in W \subseteq H(22,56)$  be as above with  $C'$  smooth and connected. If  $c_1$  is chosen among  $1 \leq c_1 \leq 9$ , then  $C' \subseteq \mathbb{P}$  defines a stable reflexive sheaf  $\underline{F}'$  and in fact a vector bundle if  $c_1 = 9$  by the usual correspondence. Using (1.1) and (2.1) we find that  $\underline{F}'$  belongs to a non-reduced component of  $M(c_1, c_2, c_3)$  for the choices  $1 \leq c_1 \leq 2$  or  $c_1 = 6$ . In particular there exists a non-reduced component of  $M(6, 22, 66) \simeq M(0, 13, 66)$ . Moreover we obtain precisely one non-reduced component in ~~this way~~  <sup>$M(6, 22, 66)$</sup>  if we make use of the discussion after (2.3). If  $c_1 = 9$ , we find a reflexive sheaf  $\underline{F}' \in M(9, 22, 0)$ , and the normalized one is  $\underline{F}'(-5) \in M(-1, 2, 0)$ , but we can not conclude that  $M(-1, 2, 0)$  is non-reduced, even though  $H(22, 56)$  is, because the condition  $H^1(\underline{I}_{C'}(c_1-4)) = 0$  of (2.1.ii) is not satisfied. In fact one knows that  $M(-1, 2, 0)$  is a smooth scheme. See [H,S] or [S,M,S].

As a starting point of these final considerations, we will suppose as known that there is an open smooth connected subscheme  $U_M \subseteq M(-1, 2, 0)$  of stable reflexive sheaves  $\underline{F}$  for which there exists a global section  $s \in H^0(\underline{F}(2))$  whose corresponding scheme of zero's  $C' = (s)_0$  is a disjoint union of two coniques. Moreover  $\dim U_M = 11$ . In fact [H,S] proves even more. We then have an exact sequence

$$0 \rightarrow \mathcal{O}_{\mathbb{P}} \rightarrow \underline{F}(2) \rightarrow \underline{I}_{C'}(3) \rightarrow 0$$

for  $\underline{F} \in U_M$ , and since the dimension of the cohomology groups  $H^i(\underline{I}_{C'}(v))$  is easily found in case  $C'$  consists of two disjoint

coniques, we get

$$h^0(\underline{F}(1)) = h^0(\underline{I}_C(2)) = 1$$

and

$$h^1(\underline{F}(v)) = h^1(\underline{I}_C(v+1)) = \begin{cases} 1 & \text{for } v = -1, 1 \\ 2 & \text{for } v = 0 \\ 0 & \text{for } v \notin \{-1, 0, 1\}. \end{cases}$$

By  $\dim U_M = 11$ ,  $\text{Ext}_{\mathbb{P}}^2(\underline{F}, \underline{F}) = 0$ . (The reader who is more familiar with the Hilbert scheme may prove our assumptions on  $U_M$  by first proving that there is an open smooth connected subscheme  $U \subseteq H(4, -1)$  of disjoint coniques  $C'$  and that  $\dim U = 16$ . This is in fact a very special case of [K1, (3.1.10 i)]. See also [K1, (3.1.4) and (2.3.18)]. With  $c_1 = 3$ , we have  $H^1(\underline{I}_C(c_1)) = H^1(\underline{I}_C(c_1-4)) = 0$ , and by the discussion after (2.3), there exists an open smooth connected subscheme of  $M(3, 4, 0) \xrightarrow{\sim} M(-1, 2, 0)$  defined by  $U_M = i(p(q^{-1}(U)))$ . Moreover  $\dim U_M = 11$  because  $\dim U_M + h^0(\underline{F}(2)) = \dim U + h^0(\omega_C(4-c_1))$ ).

Fix an integer  $v \geq 1$ , and let  $U(v)$  be the subset of  $H(d, g)$  obtained by varying  $\underline{F} \in U_M \subseteq M(-1, 2, 0)$  and by varying the sections  $s \in H^0(\underline{F}(v))$  so that  $C = (s)_0$  is a curve, i.e. let  $U(v) = q(p^{-1}(U_M))$  and regard  $U_M$  as a subscheme of  $M(c_1, c_2, 0)$  with

$$c_1 = 2v-1, \quad c_2 = 2-v+v^2, \quad d = c_2 \quad \text{and} \quad g = 1 + \frac{1}{2}c_2(c_1-4).$$

Recall that  $p$  and  $q$  are projection morphisms

$$\begin{array}{ccc} D & \xrightarrow{q} & H(d, g) \\ \downarrow p & & \\ M(c_1, c_2, 0) & & \end{array}$$

For  $(C \subseteq \mathbb{P}) \in U(v)$ , there is an exact sequence

$$0 \rightarrow 0_{\mathbb{P}} \rightarrow \underline{F}(v) \rightarrow \underline{I}_C(2v-1) \rightarrow 0$$

some  $\underline{F}(v) \in U_M$ . Now (1.1.ii) and (2.1ii) apply for  $v = 2$  and all  $v \geq 6$ , and it follows that  $H(d,g)$  is smooth at any  $(C \subseteq \mathbb{P})$  in the open subset  $U(v) \subseteq H(d,g)$ . Moreover by the irreducibility of  $p$ ,  $U(v)$  is an open smooth connected subscheme of  $H(d,g)$ .

Furthermore

$$\dim U(v) = 4d + \frac{1}{6}v(v-5)(2v-5) \quad \text{for } v \geq 6$$

(resp =  $4d$  for  $v = 2$ ) which asymptotically is  $\sim 4d + \frac{1}{3}d^{3/2}$  for  $v \gg 0$ . To find the dimension of  $U(v)$ , we use the fact that  $p$  and  $q$  are smooth morphisms of relative dimension  $h^0(\underline{F}(v)) - 1$  and  $h^0(w_C(4-c_1)) - 1$  respectively. This gives

$$\dim U_M + h^0(\underline{F}(v)) = \dim U(v) + h^0(w_C(4-c_1))$$

for  $v = 2$  and  $v \geq 6$ , and since  $h^0(w_C(4-c_1)) = h^1(O_C(c_1-4)) = 1$  for  $v \geq 6$  (resp. = 2 for  $v = 2$ ), we get

$$\dim U(v) = 10 + h^0(\underline{F}(v)) \quad \text{for } v \geq 6$$

(resp. =  $9 + h^0(\underline{F}(v))$  for  $v = 2$ ). The reader may verify that  $h^0(\underline{F}(v)) = \chi(\underline{F}(v)) = \frac{1}{6}(v-1)(2v+3)(v+4) = 4d + \frac{1}{6}(v-5)(2v-5)v - 10$  for any  $v \geq 2$ , and the conclusion follows.

We will now discuss the cases  $3 \leq v \leq 5$  where we can not guarantee the smoothness of  $q$  since (2.1.ii) does not apply. If  $v = 5$ , then the closure of  $U(5)$  in  $H(22,56)$  makes a non-reduced component by (3.3). For  $v = 3$  or  $4$ , we claim that  $H(d,g)$  is smooth along  $U(v)$  and the codimension

$$\dim W - \dim U(\nu) = h^1(\underline{I}_C(c_1-4)) = h^1(\underline{F}(-4))$$

where  $W$  is the irreducible component of  $H(d, g)$  which contains  $U(\nu)$ . To see this it suffices to prove  $H^1(\underline{N}_C) = 0$  and  $\text{Ext}^2(\underline{I}_C(c_1), \underline{F}(\nu)) = 0$  for any  $(C \subseteq \mathbb{P}) \in U(\nu)$  because these conditions imply that the scheme  $D$  and  $H(d, g)$  are non-singular at any  $(C, \xi)$  with  $\xi \in H^0(\omega_C(4-c_1))$  and  $(C \subseteq \mathbb{P}) \in H(d, g)$  respectively. See (1.1i). Moreover if these "obstruction groups" vanish, we find

$$\begin{aligned} \dim W - \dim U(\nu) &= \dim W - \dim q^{-1}(U(\nu)) = h^0(\underline{N}_C) - \dim \text{Ext}^1(\underline{I}_C(c_1), \underline{F}(\nu)) \\ &= h^1(\underline{I}_C(c_1-4)) \end{aligned}$$

where  $\dim U(\nu) = \dim q^{-1}(U(\nu))$  because of  $h^0(\omega_C(4-c_1)) = 1$ , and where the equality to the right follows from the long exact sequence of (2.2). Now to prove  $\text{Ext}^2(\underline{I}_C(c_1), \underline{F}(\nu)) = 0$  we use the long exact sequence (\*) in the proof of (1.1. i) combined with  $H^1(\underline{F}(\nu)) = 0$  and  $\text{Ext}^2(\underline{F}, \underline{F}) = 0$ , and to prove  $H^1(\underline{N}_C) = 0$  we use the long exact sequence of (2.2) combined with  $\text{Ext}^2(\underline{I}_C(c_1), \underline{F}(\nu)) = 0$  and  $\text{Ext}^3(\underline{I}_C(c_1), \mathcal{O}_{\mathbb{P}}) \simeq H^0(\underline{I}_C(c_1-4))^{\vee} = H^0(\underline{F}(\nu-4))^{\vee} = 0$  for  $\nu = 3$  or  $\nu = 4$ , and we are done.

Computing numbers, we find for  $\nu = 3$  that  $U(3)$  is a locally closed subset of  $H(8, 5)$  of codimension 1, and any smooth connected curve  $(C \subseteq \mathbb{P}) \in U(3)$  is a canonical curve, i.e.  $\omega_C \simeq \mathcal{O}_C(1)$ . For  $\nu = 4$ ,  $U(4)$  is of codimension 2 in  $H(14, 22)$  and  $\omega_C \simeq \mathcal{O}_C(\frac{2}{3})$  for any  $(C \subseteq \mathbb{P}) \in U(4)$ .

Bibliography.

- [EGA] Grothendieck, A. and Dieudonné, J.: *Eléments de Géométrie Algébriques*. Publ. Math. IHES 4 (1960), 8 (1961), 11 (1961), 17 (1963), 20 (1964), 24 (1965), 28 (1966) and 32 (1967).
- [E,S] Ellingsrud, G. and Strømme, S.A.: On the moduli space for stable rank-2 vector bundles on  $\mathbb{P}^2$ . Preprint, Univ. of Oslo, No 6 (1979).
- [G] Gieseker, D.: On the moduli of vector bundles on an algebraic surface. *Ann. Math.* 106, 45-60 (1977).
- [H1] Hartshorne, R.: *Algebraic Geometry*. Graduate Texts in Math., Vol. 52, Springer-Verlag, New York (1977).
- [H2] Hartshorne, R.: Stable vector bundles of rank 2 on  $\mathbb{P}^3$ . *Math. Ann.* 238, 229-280 (1978).
- [H3] Hartshorne, R.: Stable Reflexive Sheaves. *Math. Ann.* 254, 121-176 (1980).
- [H,S] Hartshorne, R. and Sols, I.: Stable rank 2 vector bundles on  $\mathbb{P}^3$  with  $c_1 = -1$ ,  $c_2 = 2$ . Preprint (1980).
- [K1] Kleppe, J.O.: The Hilbert-flag scheme, its properties and its connection with the Hilbert scheme. Applications to curves in 3-space. Preprint, Univ. of Oslo, No 5 (1981)
- [L1] Laudal, O.A.: Formal Moduli of Algebraic Structures. Springer Lecture Notes nr. 754 (1979).
- [L2] Laudal, O.A.: A generalized tri-secant lemma. Proc. of t. Tromsø alg. geom. Conference 1977. Springer Lecture Notes nr. 687, 112-149 (1978).
- [M1] Mumford, D.: Further pathologies in algebraic geometry. *Amer. J. Math.* 84, 642-647 (1962).

- [M2] Mumford, D.: Geometric Invariant Theory, Springer Verlag Berlin-Heidelberg-New York (1965).
- [Ma 1] Maruyama, M.: Moduli of stable sheaves I, J. Math., Kyoto Univ. 17, 91-166 (1977).
- [Ma 2] Maruyama, M.: Moduli of stable sheaves II, J. Math., Kyoto Univ. 18, 557-614 (1978).
- [S,P] Szpiro, L. and Peskine, C.: Liaison des variétés algébriques. Invent. math. 26, 271-302 (1974).
- [S,M,S] Strømme, S.A., Mesequer, J. and Sols, I.: Compactification of a Family of Vector Bundles on  $\mathbb{P}^3$ . Proc. of t. 18th Scandinavian Congress of Math. (1980).